



The area of a self-similar fragmentation

Jean Bertoin

Institut für Mathematik, Universität Zürich
 Winterthurerstrasse 190,
 CH-8057 Zürich, Switzerland.

E-mail address: jean.bertoin@math.uzh.ch

URL: <http://www.math.uzh.ch/index.php?professur&key1=6119>

Abstract. We consider the area $A = \int_0^\infty (\sum_{i=1}^\infty X_i(t)) dt$ of a self-similar fragmentation process $\mathbf{X} = (\mathbf{X}(t), t \geq 0)$ with negative index. We characterize the law of A by an integro-differential equation. The latter may be viewed as the infinitesimal version of a recursive distribution equation that arises naturally in this setting. In the case of binary splitting, this yields a recursive formula for the entire moments of A which generalizes known results for the area of the Brownian excursion.

1. Introduction

The distribution of the area $A_{\text{Exc}} = \int_0^1 e_s ds$ of a standard Brownian excursion ($e_s, 0 \leq s \leq 1$) appears in a variety of settings, including random graphs cf. [Spencer \(1997\)](#); [Takács \(1991a\)](#), random trees and branching processes cf. [Takács \(1991b, 1994\)](#), order statistics cf. [Takács \(1992\)](#), hashing with linear probing cf. [Flajolet et al. \(1998\)](#), ..., not to mention of course the study of Brownian motion for its own interest cf. [Louchard \(1984a,b\)](#). The entire moments $\mathbb{E}(A_{\text{Exc}}^k)$ have a special importance, as they are related, for instance, to asymptotics as $n \rightarrow \infty$ for the number of connected graphs with n labelled vertices and $n + k - 1$ edges, see [Spencer \(1997\)](#) and the survey by [Janson \(2007\)](#). We refer to [Perman and Wellner \(1996\)](#); [Janson \(2007\)](#) and references therein for a detailed presentation and reviews of known results on this topic.

The starting point of this work lies in the observation that one can express the area in the form

$$A_{\text{Exc}} = \int_0^\infty |\theta(t)| dt$$

where $|\theta(t)|$ denotes the Lebesgue measure of the random open set $\theta(t) = \{s \in [0, 1] : e_s > t\}$. The point is that the process $\theta = (\theta(t), t \geq 0)$ is a self-similar

Received by the editors January 20, 2011; accepted November 23, 2011.

2000 Mathematics Subject Classification. 60G18; 60J80.

Key words and phrases. Self-similar fragmentation, area, recursive distributional equation.

Research supported by ANR-08-BLAN-0220-01.

interval-fragmentation in the sense of Bertoin (2002, 2006). One can derive an integro-differential equation for the distribution of A_{Exc} from the branching and self-similarity properties of θ . In particular this yields recursive formulas for the entire moments of A_{Exc} that have been obtained in the literature by analytic techniques based on the Feynman-Kac formula or the analysis of continued fractions and of singularities of the generating functions of discrete approximations.

The same approach applies more generally to self-similar fragmentation processes, a class of Feller processes with values in the space of mass-partitions

$$\mathcal{P}_m = \{\mathbf{x} = (x_1, x_2, \dots) : x_1 \geq x_2 \geq \dots \geq 0 \text{ and } \sum_{i=1}^{\infty} x_i \leq 1\}.$$

Specifically, a self-similar fragmentation $\mathbf{X} = (\mathbf{X}(t), t \geq 0)$ fulfills the following two fundamental properties. For $x \in [0, 1]$, let us denote by \mathbb{P}_x the law of the version of \mathbf{X} started from a single fragment of mass x , i.e. $\mathbf{X}(0) = (x, 0, \dots)$. First, the self-similarity means that there exists an index $\alpha \in \mathbb{R}$ such that for every $x \in (0, 1]$ the distribution of the rescaled process $(x\mathbf{X}(x^\alpha t), t \geq 0)$ under \mathbb{P}_1 is \mathbb{P}_x . Second, the process \mathbf{X} satisfies the branching property, in the sense that for every mass-partition $\mathbf{x} = (x_1, x_2, \dots)$, if $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots$ are independent fragmentations with respective laws $\mathbb{P}_{x_1}, \mathbb{P}_{x_2}, \dots$, then the process resulting from the decreasing rearrangement of all the fragments of $\mathbf{X}^{(1)}(t), \mathbf{X}^{(2)}(t), \dots$ is a version of $\mathbf{X}(t)$ started from $\mathbf{X}(0) = \mathbf{x}$.

We assume that the index of self-similarity α is negative, which implies that small fragments split faster than the large ones. A well-known consequence is that the process of the total mass $t \mapsto \sum_{i=1}^{\infty} X_i(t)$ decreases and reaches 0 in finite time a.s.; in other words the entire mass is eventually ground down to dust. This has been observed first by Filippov (1961), see also Proposition 2(i) in Bertoin (2003). We may thus define the area

$$A = \int_0^\infty \left(\sum_{i=1}^{\infty} X_i(t) \right) dt$$

which is the main object of interest in this work. The denomination *area* is better understood if we remember that a fragmentation admits an interval representation; cf. Section 3.2 in Bertoin (2002). There exists a nested right-continuous family $(\theta(t), t \geq 0)$ of open subsets of the unit interval such that for every $t \geq 0$, the sequence of the lengths of the interval components of $\theta(t)$ listed in the decreasing order is precisely $\mathbf{X}(t) = (X_1(t), \dots)$. If we define a lower semi-continuous path $F : [0, 1] \rightarrow \mathbb{R}_+$ by

$$F(u) = \sup\{t \geq 0 : u \in \theta(t)\}, \quad u \in [0, 1],$$

then $\theta(t) = \{u : F(u) > t\}$, and since $\sum_{i=1}^{\infty} X_i(t) = |\theta(t)|$, we can express A in the form $A = \int_0^1 F(u) du$. Of course F is the Brownian excursion in the situation discussed at the beginning of this introduction.

The area A has another natural interpretation in terms of continuous random trees. Indeed, Haas and Miermont (2004) obtained a representation of self-similar fragmentations in terms of some random rooted real tree (\mathbf{T}, μ) (where μ denotes a probability measure on the metric space \mathbf{T}) which enjoys self-similarity and branching properties. More precisely, for every $t \geq 0$, $\mathbf{X}(t) = (X_1(t), \dots)$ can be viewed as the ranked sequence of the masses of connected components of $\mathbf{T}(t)$, the subset of the points in \mathbf{T} at distance at least t from the root. Thus in this setting,

$\mu(\mathbf{T}(t)) = \sum_{i=1}^{\infty} X_i(t)$ and

$$A = \int_0^{\infty} \mu(\mathbf{T}(t)) dt.$$

In other words, A represents the *average height* in \mathbf{T} , i.e. the average distance of points to the root, where averaging is taken with respect to the probability measure μ on \mathbf{T} . Beware that this differs from the *typical height* H , that is the distance from the root of a point chosen uniformly at random according to μ . More precisely, A is the conditional expectation of H given (\mathbf{T}, μ) . We also refer to [Takács \(1994\)](#) for results on this quantity in the framework of certain discrete random trees.

In the next section, we will present our main result which determines the law of A as the unique solution to an intro-differential equation expressed in terms of the characteristics of \mathbf{X} . In the case of binary dislocations, this enables us to derive explicit recursive formulas for the moments of A . We recover in particular identities due to [Takács \(1991a\)](#) for the moments of the area of the Brownian excursion. Section 3 is devoted to the proof of this integro-differential equation. We shall start by establishing a priori bounds for the moments of A . Then we proceed with the simpler case when the dislocation measure is finite, and derive the equation from a recursive distribution equation which is naturally induced by the dynamics of fragmentation. The general case when the dislocation measure is infinite is then deduced by approximation. This relies on a weak limit theorem for the area of fragmentation processes. The proof of the latter is somewhat technical and will be postponed to the final subsection.

2. Main results

We denote by ν the dislocation measure of \mathbf{X} , so ν is a sigma-finite measure on the space of mass-partitions \mathcal{P}_m with no atom at the trivial mass-partition $(1, 0, \dots)$ and fulfills the integral condition

$$\int_{\mathcal{P}_m} (1 - x_1) \nu(d\mathbf{x}) < \infty. \quad (2.1)$$

We implicitly exclude the degenerate case when $\nu \equiv 0$ and further assume absence of erosion. Roughly speaking, this means that \mathbf{X} is a purely discontinuous process that only evolves by sudden dislocations whose rates are determined by ν and the index of self-similarity α . We refer to Chapter 3 of [Bertoin \(2006\)](#) or [Bertoin \(2002\)](#) for background.

For an arbitrary mass partition \mathbf{x} , we denote by

$$\eta_{\mathbf{x}}(da) = \mathbb{P}_{\mathbf{x}}(A \in da), \quad a \geq 0,$$

the law of the area under the probability measure $\mathbb{P}_{\mathbf{x}}$ for which the fragmentation \mathbf{X} starts from $\mathbf{X}(0) = \mathbf{x}$. For the sake of simplicity, we will work from now on under the law $\mathbb{P} = \mathbb{P}_1$, i.e. when the fragmentation starts from a single fragment of mass 1, and write then $\eta = \eta_1$. This induces no loss of generality since, combining self-similarity and the branching property, we get that for every mass-partition $\mathbf{x} = (x_1, x_2, \dots)$, there is the identity

$$\eta_{\mathbf{x}}(da) = \mathbb{P} \left(\sum_{i=1}^{\infty} x_i^{1-\alpha} A_i \in da \right) \quad (2.2)$$

where $(A_i)_{i \in \mathbb{N}}$ is a family of i.i.d. copies of A . Note that when \mathbf{x} has only finitely many non-zero terms, say x_1, \dots, x_n , then $\eta_{\mathbf{x}}$ can be expressed as a convolution product $\eta_{\mathbf{x}} = \eta_{x_1} * \dots * \eta_{x_n}$ where η_y stands for the image of η by the dilation $a \mapsto y^{1-\alpha}a$. Finally, we let $\langle \mu, f \rangle = \int f d\mu$ denote the integral of some function f with respect to a measure μ when the integral makes sense. We are now able to state our main result.

Theorem 2.1. *Let \mathbf{X} be a self-similar fragmentation with index $\alpha < 0$, dislocation measure ν and without erosion. Then for every \mathcal{C}^1 function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ such that $f'(y) = O(y^p)$ as $y \rightarrow \infty$ for some $p > 0$, the law η of A solves*

$$\langle \eta, f' \rangle = \int_{\mathcal{P}_m} \nu(d\mathbf{x}) (\langle \eta, f \rangle - \langle \eta_{\mathbf{x}}, f \rangle), \quad (2.3)$$

where the quantities above are finite. Further (2.3) characterizes η .

If we introduce the concave increasing function $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ by

$$\Phi(q) = \int_{\mathcal{P}_m} \left(1 - \sum_{j=1}^{\infty} x_j^{1+q} \right) \nu(d\mathbf{x}), \quad (2.4)$$

then we immediately see from Theorem 2.1 that the first moment of A is given simply by

$$\mathbb{E}(A) = \langle \eta, \text{Id} \rangle = 1/\Phi(-\alpha).$$

This identity can also be established directly; see the forthcoming Lemma 3.1.

More generally, we shall now derive from Theorem 2.1 a recursive formula for the entire moments of A . For the sake of simplicity, we shall focus on the special case of binary dislocations, although more general situations could be dealt with at the price of heavier notation. This means that we assume that the dislocation measure ν has support in the subset of binary mass-partitions $\{\mathbf{x} = (x, 1-x, 0, \dots), x \in [1/2, 1)\}$. By a slight abuse, we shall then identify the dislocation measure ν with its image by the map $\mathbf{x} \rightarrow x_1$, i.e. we view ν as a measure on $[1/2, 1)$. Specializing Theorem 2.1 to $f(x) = x^k$ and applying the binomial formula, we immediately obtain:

Corollary 2.2. *Let \mathbf{X} be a self-similar fragmentation with index $\alpha < 0$, binary dislocation measure ν and without erosion. For every integer $k \geq 0$, let $M_k = \mathbb{E}(A^k)$ denote the k -th moment of the area; in particular $M_0 = 1$. Then there is the identity*

$$a_k M_k = k M_{k-1} + \sum_{j=1}^{k-1} a_{j,k} M_j M_{k-j}, \quad k \geq 1,$$

with

$$a_k = \int_{[1/2, 1)} (1 - x^{k(1-\alpha)} - (1-x)^{k(1-\alpha)}) \nu(dx) = \Phi(k(1-\alpha) - 1),$$

and

$$a_{j,k} = \binom{k}{j} \int_{[1/2, 1)} x^{j(1-\alpha)} (1-x)^{(k-j)(1-\alpha)} \nu(dx).$$

We stress that (2.1) ensures the finiteness of a_k and $a_{j,k}$.

We now discuss some examples, starting with the case of the Brownian fragmentation $A = A_{\text{Exc}}$ which has motivated this work. The Brownian fragmentation has

self-similarity index $\alpha = -1/2$, no erosion, and its dislocation measure is binary and specified by

$$\nu(dx) = \frac{2}{\sqrt{2\pi x^3(1-x)^3}} dx, \quad 1/2 \leq x < 1;$$

see [Bertoin \(2002\)](#) on its pages 339-340. One gets by symmetry

$$a_k = \int_0^1 \frac{1 - x^{3k/2} - (1-x)^{3k/2}}{\sqrt{2\pi x^3(1-x)^3}} dx = 2^{3/2} \frac{\Gamma((3k-1)/2)}{\Gamma(3k/2-1)}$$

and

$$\begin{aligned} a_{j,k} &= \binom{k}{j} \int_0^1 \frac{x^{3j/2}(1-x)^{3(k-j)/2}}{\sqrt{2\pi x^3(1-x)^3}} dx \\ &= \frac{k! \Gamma((3j-1)/2) \Gamma((3(k-j)-1)/2)}{\sqrt{2\pi} j! (k-j)! \Gamma(3k/2-1)} \end{aligned}$$

Following [Takács \(1991a\)](#), if we set

$$M_k = \frac{4\sqrt{\pi} 2^{-k/2} k!}{\Gamma((3k-1)/2)} K_k,$$

then after some cancellations, [Corollary 2.2](#) reduces to

$$K_k = (3k/4 - 1)K_{k-1} + \sum_{j=1}^{k-1} K_j K_{k-j}$$

with $K_0 = -1/2$. This is the recursive equation found by [Takács](#), which in turn yields a Riccati type ODE by considering the exponential generating function of the K_k ; see [Flajolet et al. \(1998\)](#). We mention the existence of other recursive formulas for the moments of A_{Exc} , see in particular [Louchard \(1984a\)](#) and the discussion in [Janson \(2007\)](#).

Similar calculations apply when more generally the dislocation measure is of beta-type, i.e. is binary with

$$\nu(dx) = cx^\beta(1-x)^\beta dx, \quad 1/2 < x < 1$$

for some $-2 < \beta < -1$. These beta-splitting measures have appeared in works of [Aldous \(1996\)](#) on cladograms; see also Section 5.1 in [Haas et al. \(2008\)](#). One obtains

$$\begin{aligned} & \frac{2}{c} a_k \\ &= \int_0^1 \left(1 - x^{k(1-\alpha)} - (1-x)^{k(1-\alpha)}\right) x^\beta (1-x)^\beta dx \\ &= \text{B}(\beta+1, \beta+1) - 2\text{B}(\beta+1+k(1-\alpha), \beta+1) \\ &= \frac{2(2\beta+3)}{\beta+1} \text{B}(\beta+2, \beta+2) - 2 \frac{2\beta+2+k(1-\alpha)}{\beta+1} \text{B}(\beta+1+k(1-\alpha), \beta+2) \end{aligned}$$

where $\text{B}(a, b) = \Gamma(a)\Gamma(b)/\Gamma(a+b)$ is the beta function, and

$$\begin{aligned} \frac{2}{c} a_{j,k} &= \binom{k}{j} \int_0^1 x^{\beta+j(1-\alpha)} (1-x)^{\beta+(k-j)(1-\alpha)} dx \\ &= \binom{k}{j} \text{B}(\beta+j(1-\alpha)+1, \beta+(k-j)(1-\alpha)+1). \end{aligned}$$

Finally, note that we can also deal with linear combinations of the beta dislocation measures, which covers for instance the case of Ford's alpha model; see Section 5.2 in [Haas et al. \(2008\)](#).

3. Proof of Theorem 2.1

This section is devoted to the proof of Theorem 2.1; it relies in four main steps. In the first sub-section, we establish a priori bounds for the moments of the area, relying on known properties of the so-called tagged fragment. In the second sub-section, we prove the equation (2.3) in the special case when the dislocation measure is finite. In the third sub-section, we provide the proof of Theorem 2.1 by approximation, taking for granted a weak convergence result for the area that will be established in the final sub-section.

3.1. *Bounds for the moments of the area.* The purpose of this subsection is to establish some a priori bounds on the moments $M_k = \mathbb{E}(A^k)$ of the area. Recall the notation (2.4) and that $\alpha < 0$.

Lemma 3.1. *We have*

$$M_1 = 1/\Phi(-\alpha)$$

and for $k \geq 1$

$$M_k \leq \frac{k!}{\Phi(-\alpha) \cdots \Phi(-k\alpha)}.$$

As a consequence $\mathbb{E}(\exp(cA)) < \infty$ whenever $c < \Phi(\infty)$, and in particular the law η of A is determined by its entire moments.

Proof: It is convenient to work in the framework of interval-fragmentation, i.e. when the fragmentation \mathbf{X} describes the ranked sequence of the lengths of the interval components of a family $(\theta(t), t \geq 0)$ of nested open subsets of the unit interval. Recall from the introduction that if we set

$$F(u) = \sup\{t \geq 0 : u \in \theta(t)\}, \quad u \in [0, 1],$$

then $A = \int_0^1 F(u)du$, and then for every integer $k \geq 1$, there is the identity

$$A^k = \int_0^1 du_1 \cdots \int_0^1 du_k F(u_1) \cdots F(u_k).$$

In this setting, we have

$$\mathbb{E}(A^k) = \mathbb{E}(F(U_1) \cdots F(U_k))$$

where U_1, \dots, U_k are i.i.d. uniform variables on $[0, 1]$. This yields

$$M_1 = \mathbb{E}(F(U)) \quad \text{and} \quad M_k \leq \mathbb{E}(F(U))^k$$

where U has the uniform distribution on $[0, 1]$.

The variable $F(U)$ should be viewed as the lifetime of the tagged-fragment, i.e. it is the first instant t when the size $\chi(t)$ of the interval component of $\theta(t)$ that contains the randomly tagged point U reaches the absorbing state 0. This variable has the distribution of an exponential functional,

$$F(U) \stackrel{(\text{law})}{=} I = \int_0^\infty \exp(\alpha \xi_t) dt$$

where $\xi = (\xi_t, t \geq 0)$ is a subordinator with Laplace exponent Φ ; see Corollary 2 in Bertoin (2002). Since it is well-known that

$$\mathbb{E}(I^k) = \frac{k!}{\Phi(-\alpha) \cdots \Phi(-k\alpha)}$$

(cf. for instance Theorem 2 in Bertoin and Yor, 2005), the first two claims are proved, and the last ones follow immediately. Indeed, since the function Φ increases, we have for every $c < b < \Phi(\infty)$ that $\Phi(-\alpha) \cdots \Phi(-k\alpha) \geq b^k$ provided that k is sufficiently large. In particular, we deduce from the bounds for M_k that the series $\sum_{k=0}^{\infty} \frac{c^k M_k}{k!}$ converges, that is $\mathbb{E}(\exp(cA)) < \infty$. It is well-known that this guaranties that the moment problem for the law of A is determined. \square

3.2. The case with finite dislocation rates. In this subsection, we assume that the fragmentation process has a finite dislocation measure, i.e. $\nu(\mathcal{P}_m) \in (0, \infty)$. This means that under \mathbb{P} , the process \mathbf{X} stays in state $(1, 0, \dots)$ during an exponential time T with parameter $\nu(\mathcal{P}_m)$, and then, independently of the waiting time T , jumps at some random mass partition $\mathbf{X}(T)$ whose distribution is given by the normalized dislocation measure $\nu/\nu(\mathcal{P}_m)$. We stress that the jump times of \mathbf{X} may nonetheless accumulate right after T ; in particular then \mathbf{X} is not a continuous-time Markov chain. Indeed, the first dislocation may produce fragments of arbitrarily small sizes, which then split again almost instantaneously by self-similarity.

The proof of the following weaker version of Theorem 2.1 in this setting is straightforward.

Proposition 3.2. *Assume that \mathbf{X} has no erosion and finite dislocation measure ν . Then for every \mathcal{C}^1 function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ such that the derivative f' has a finite limit at ∞ , we have*

$$\langle \eta, f' \rangle = \int_{\mathcal{P}_m} \nu(d\mathbf{x}) (\langle \eta, f \rangle - \langle \eta_{\mathbf{x}}, f \rangle).$$

Proof: An application of the strong Markov property at the first dislocation time T and (2.2) yields the recursive distributional equation (see the survey by Aldous and Bandyopadhyay (2005) for much more this topic)

$$A = T + \sum_{i=1}^{\infty} X_i(T)^{1-\alpha} A_i$$

where $(A_i)_{i \in \mathbb{N}}$ is a sequence of i.i.d. copies of A which is further independent of $\mathbf{X}(T)$. This entails

$$\ell(q) = \frac{1}{\nu(\mathcal{P}_m) + q} \int_{\mathcal{P}_m} \nu(d\mathbf{x}) \prod_{i=1}^{\infty} \ell(x_i^{1-\alpha} q), \quad q \geq 0$$

where $\ell(q) = \mathbb{E}(\exp(-qA))$ is Laplace transform of the area. By rearrangement, we arrive at

$$-q\ell(q) = \int_{\mathcal{P}_m} \nu(d\mathbf{x}) \left(\ell(q) - \prod_{i=1}^{\infty} \ell(x_i^{1-\alpha} q) \right),$$

which is the equation in the statement specified for the exponential function $f(a) = e^{-qa}$.

We shall now extend this to more general functions by an application of the Stone-Weierstrass theorem. Recall that the latter ensures that linear combinations

of exponential functions $a \rightarrow e^{-qa}$ with $q \geq 0$ are everywhere dense in the space $\mathcal{C}([0, \infty], \mathbb{R})$ of continuous functions on the half-line with a finite limit at infinity, endowed with the uniform norm $\|g\|_\infty = \sup_{x \geq 0} |g(x)|$. Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that its derivative f' belongs to $\mathcal{C}([0, \infty], \mathbb{R})$. So there exists a sequence $(f_n)_{n \in \mathbb{N}}$ of linear combinations of exponential functions with $f_n(0) = f(0)$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} \|f'_n - f'\|_\infty = 0$. In particular f_n converges to f uniformly on compact sets and there exists some constant $c > 0$ such that $|f_n(x)| \leq c(1+x)$ and $|f(x)| \leq c(1+x)$ for all $x \geq 0$.

Therefore we have

$$\lim_{n \rightarrow \infty} \langle \eta, f'_n \rangle = \langle \eta, f' \rangle$$

and for every $a > 0$ and every mass-partition $\mathbf{x} \in \mathcal{P}_m$ (including possibly $\mathbf{x} = (1, 0, \dots)$)

$$\lim_{n \rightarrow \infty} \langle \eta_{\mathbf{x}}, \mathbf{1}_{[0,a]} f_n \rangle = \langle \eta_{\mathbf{x}}, \mathbf{1}_{[0,a]} f \rangle.$$

Recall that ν is assumed to be finite in this section; since the functions $\mathbf{1}_{[0,a]} f_n$ are bounded uniformly in $n \in \mathbb{N}$ and the $\eta_{\mathbf{x}}$ are probability measures, we may apply dominated convergence and obtain

$$\lim_{n \rightarrow \infty} \int_{\mathcal{P}_m} \nu(d\mathbf{x}) (\langle \eta, \mathbf{1}_{[0,a]} f_n \rangle - \langle \eta_{\mathbf{x}}, \mathbf{1}_{[0,a]} f_n \rangle) = \int_{\mathcal{P}_m} \nu(d\mathbf{x}) (\langle \eta, \mathbf{1}_{[0,a]} f \rangle - \langle \eta_{\mathbf{x}}, \mathbf{1}_{[0,a]} f \rangle).$$

On the other hand, since $|f_n(x)| \leq c(1+x)$, we have for every mass-partition $\mathbf{x} \in \mathcal{P}_m$

$$|\langle \eta_{\mathbf{x}}, \mathbf{1}_{(a,\infty)} f_n \rangle| \leq c \mathbb{E}_{\mathbf{x}}(1+A, A > a),$$

and the same upper bound holds with f replacing f_n in the left-hand side. Note that

$$\mathbb{E}_{\mathbf{x}}(A) = \sum_1^\infty x_i^{1-\alpha} \mathbb{E}(A) \leq \mathbb{E}(A) = M_1,$$

where $\mathbf{x} = (x_1, \dots)$ denotes an arbitrary mass-partition, so Markov inequality yields the uniform bound

$$\mathbb{P}_{\mathbf{x}}(A > a) \leq a^{-1} M_1.$$

Moreover,

$$\mathbb{E}_{\mathbf{x}}(A^2) = \sum_{i=1}^\infty \sum_{j=1}^\infty x_i^{1-\alpha} x_j^{1-\alpha} \mathbb{E}(A_i A_j) \leq \mathbb{E}(A^2) = M_2$$

where $(A_i)_{i \in \mathbb{N}}$ is a sequence of i.i.d. copies of A , so by Hölder inequality

$$\mathbb{E}_{\mathbf{x}}(A, A > a) \leq \sqrt{\mathbb{P}_{\mathbf{x}}(A > a) \mathbb{E}_{\mathbf{x}}(A^2)} \leq \sqrt{a^{-1} M_1 M_2}.$$

Putting the pieces together, we derive from the triangle inequality that

$$\left| \int_{\mathcal{P}_m} \nu(d\mathbf{x}) (\langle \eta, \mathbf{1}_{(a,\infty)} f_n \rangle - \langle \eta_{\mathbf{x}}, \mathbf{1}_{(a,\infty)} f_n \rangle) \right| \leq 2c\nu(\mathcal{P}_m) \left(a^{-1} M_1 + \sqrt{a^{-1} M_1 M_2} \right)$$

and the same inequality holds when f replaces f_n in the left-hand side. Recall from Lemma 3.1 that the moments M_1 and M_2 are finite, so the right-hand side can be made as small as we wish by choosing a sufficiently large.

We know from the first part of the proof that for the linear combinations f_n of exponential functions, there is the identity

$$\langle \eta, f'_n \rangle = \int_{\mathcal{P}_m} \nu(d\mathbf{x}) (\langle \eta, f_n \rangle - \langle \eta_{\mathbf{x}}, f_n \rangle),$$

and we have all the ingredients to take limits as $n \rightarrow \infty$. We conclude that the same formula holds with f replacing f_n . \square

Let us briefly discuss the elementary example when $\nu = \delta_{(1/2, 1/2, 0, \dots)}$. We thus start with a single fragment of unit size which splits in two fragments each of size $1/2$ after an exponential time with parameter 1, and so on. It should be plain that the area can then be expressed in the form

$$A = \mathbf{e}_{0,1} + 2^{\alpha-1}(\mathbf{e}_{1,1} + \mathbf{e}_{1,2}) + 2^{2(\alpha-1)}(\mathbf{e}_{2,1} + \mathbf{e}_{2,2} + \mathbf{e}_{2,3} + \mathbf{e}_{2,4}) + \dots,$$

where the $\mathbf{e}_{i,j}$ for $j = 1, \dots, 2^i$ and $i = 0, 1, \dots$ are i.i.d. standard exponential variables. The Laplace transform of A is thus given by

$$\ell(q) = \prod_{n=0}^{\infty} \left(\frac{1}{1 + 2^{n(\alpha-1)}q} \right)^{2^n},$$

and the equation

$$-q\ell(q) = \ell(q) - \ell(2^{\alpha-1}q)^2$$

provided by Proposition 3.2 can be checked directly.

It may be interesting to observe that the very same argument as in the proof of Proposition 3.2 can be applied when analyzing the distribution of the *total length* rather than the area of a self-similar fragmentation. Specifically, consider another self-similar fragmentation \mathbf{X}' with no erosion, finite dislocation measure ν and index of self-similarity $\alpha' = \alpha - 1$. Since $\alpha' < -1$, it is known (see Proposition 2 in Bertoin (2003) and its proof) that the variable

$$L' = \int_0^{\infty} \text{Card}\{i \in \mathbb{N} : X'_i(t) > 0\} dt$$

is finite a.s. The quantity L' should be viewed as the total length of the fragmentation tree \mathbf{T}' associated by Haas and Miermont (2004) to \mathbf{X}' . Just as in the proof of Proposition 3.2, it is easily seen that there is the identity

$$L' = T' + \sum_{i=1}^{\infty} X'_i(T')^{1-\alpha} L'_i$$

where T' denotes the first dislocation time of \mathbf{X}' and $(L'_i)_{i \in \mathbb{N}}$ is a sequence of i.i.d. copies of L' which is further independent of $\mathbf{X}'(T')$ and T' . Because the dislocation measures of \mathbf{X} and \mathbf{X}' are identical, $\mathbf{X}'(T')$ has the same law as $\mathbf{X}(T)$ and T' is also an exponential variable with parameter $\nu(\mathcal{P}_m)$. We can then deduce from the arguments similar to those in the proof of Proposition 3.2 that

$$\langle \eta', f' \rangle = \int_{\mathcal{P}_m} \nu(d\mathbf{x}) (\langle \eta', f \rangle - \langle \eta'_{\mathbf{x}}, f \rangle),$$

where $\eta'_{\mathbf{x}}$ denote the distribution of L' under $\mathbb{P}'_{\mathbf{x}}$ and $\eta' = \eta'_1$. Observe that self-similarity implies that

$$\eta'_{\mathbf{x}}(da) = \mathbb{P} \left(\sum_{i=1}^{\infty} x_i^{1-\alpha} L'_i \in da \right),$$

so we conclude from Theorem 2.1 that L' has the same law as A . We point out that this identity in distribution can also be established from Equation (10) of Bertoin (2003) as a consequence of the connexion between self-similar fragmentations with

the same dislocation measure but different indices of self-similarity, and also holds when the dislocation measure ν is infinite.

3.3. Proof of Theorem 2.1 by approximation. In this subsection, we shall derive Theorem 2.1 by approximation from the case when the dislocation measure is finite, taking for granted the weak convergence of the corresponding areas. Specifically, we introduce the finite measures

$$\nu^{(n)}(d\mathbf{x}) = \mathbf{1}_{\{1-x_1 > 1/n\}} \nu(d\mathbf{x}), \quad \mathbf{x} \in \mathcal{P}_m,$$

where n is a sufficiently large integer so that $\nu^{(n)} \not\equiv 0$. We write $A^{(n)}$ for the area of a self-similar fragmentation process with index α , dislocation measure $\nu^{(n)}$ and without erosion, and denote by $\eta^{(n)}$ the distribution of $A^{(n)}$. Recall (2.2) and set for a generic mass-partition \mathbf{x}

$$\eta_{\mathbf{x}}^{(n)}(da) = \mathbb{P} \left(\sum_{i=1}^{\infty} x_i^{1-\alpha} A_i^{(n)} \in da \right)$$

where $(A_i^{(n)} : i \in \mathbb{N})$ is a sequence of i.i.d. copies of $A^{(n)}$. The following crucial lemma will be established in the next sub-section.

Lemma 3.3. *The sequence $(\eta^{(n)}, n \in \mathbb{N})$ converges weakly to η as $n \rightarrow \infty$.*

The next step to the proof of Theorem 2.1 is the following technical result.

Lemma 3.4. (i) *Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ be continuous and bounded. Then for every $\mathbf{x} \in \mathcal{P}_m$,*

$$\lim_{n \rightarrow \infty} \langle \eta_{\mathbf{x}}^{(n)}, f \rangle = \langle \eta_{\mathbf{x}}, f \rangle.$$

(ii) *Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function with $f'(y) = O(y^p)$ as $y \rightarrow \infty$ for some $p > 0$, and set $\|f'\| = \sup\{|f'(x)|/(1+x)^p : x \geq 0\}$. There is a constant c depending only on p and the characteristics of the fragmentation such that for every mass-partition \mathbf{x} and every n*

$$|\langle \eta^{(n)}, f \rangle - \langle \eta_{\mathbf{x}}^{(n)}, f \rangle| \leq c \|f'\| (1 - x_1).$$

Proof: (i) Denote by κ the cumulant of A , i.e. $\mathbb{E}(\exp(-qA)) = \exp(-\kappa(q))$ for $q \geq 0$, and by $\kappa^{(n)}$ that of $A^{(n)}$. If $(A_i^{(n)}, i \in \mathbb{N})$ is a sequence of i.i.d. copies of $A^{(n)}$, then we have

$$\mathbb{E} \left(\exp \left(-q \sum_{i=1}^{\infty} x_i^{1-\alpha} A_i^{(n)} \right) \right) = \exp \left(- \sum_{i=1}^{\infty} \kappa^{(n)}(x_i^{1-\alpha} q) \right).$$

We know from Lemma 3.3 that $\lim_{n \rightarrow \infty} \kappa^{(n)}(x_i^{1-\alpha} q) = \kappa(x_i^{1-\alpha} q)$ for every $q \geq 0$ and $i \in \mathbb{N}$. Further, $\kappa^{(n)}$ is a concave increasing function with $\kappa^{(n)}(0) = 0$, and its derivative at 0 is given by $\mathbb{E}(A^{(n)})$. Recall also from Lemma 3.1 that $\mathbb{E}(A^{(n)}) = 1/\Phi^{(n)}(-\alpha)$ where

$$\Phi^{(n)}(q) = \int_{\mathcal{P}_m} \left(1 - \sum_{i=1}^{\infty} x_i^{1+q} \right) \nu^{(n)}(d\mathbf{x}) = \int_{\mathcal{P}_m} \mathbf{1}_{\{1-x_1 > 1/n\}} \left(1 - \sum_{i=1}^{\infty} x_i^{1+q} \right) \nu(d\mathbf{x}).$$

Plainly the sequence $\Phi^{(n)}(-\alpha)$ increases and there are thus the bounds

$$\kappa^{(n)}(x_i^{1-\alpha} q) \leq x_i^{1-\alpha} q / \Phi^{(n)}(-\alpha) \leq c x_i^{1-\alpha} q.$$

Since $\sum_{i=1}^{\infty} x_i^{1-\alpha} \leq 1$, we conclude by dominated convergence that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^{\infty} \kappa^{(n)}(x_i^{1-\alpha} q) = \sum_{i=1}^{\infty} \kappa(x_i^{1-\alpha} q).$$

By Laplace inversion, this yields our claim.

(ii) Recall that $A_1^{(n)}, \dots$ are i.i.d. copies of $A^{(n)}$, and set $S_{\mathbf{x}}^{(n)} = \sum_{i=2}^{\infty} x_i^{1-\alpha} A_i^{(n)}$ where $\mathbf{x} = (x_1, \dots)$ is a generic mass-partition. We have

$$\begin{aligned} |\langle \eta^{(n)}, f \rangle - \langle \eta_{\mathbf{x}}^{(n)}, f \rangle| &\leq \mathbb{E} \left(\left| f(A_1^{(n)}) - f(x_1^{1-\alpha} A_1^{(n)} + S_{\mathbf{x}}^{(n)}) \right| \right) \\ &\leq \|f'\| \mathbb{E} \left((1 + A_1^{(n)} + S_{\mathbf{x}}^{(n)})^p \left((1 - x_1^{1-\alpha}) A_1^{(n)} + S_{\mathbf{x}}^{(n)} \right) \right). \end{aligned}$$

Recall that $\sum_1^{\infty} x_i \leq 1$. Applying Rosenthal's inequality for the moments of sum of independent nonnegative variables and then Jensen's inequality, we obtain

$$\mathbb{E} \left((1 + A_1^{(n)} + S_{\mathbf{x}}^{(n)})^{2p} \right) \leq c(2p)(1 + M_{2p}^{(n)})$$

and

$$\mathbb{E} \left(\left((1 - x_1^{1-\alpha}) A_1^{(n)} + S_{\mathbf{x}}^{(n)} \right)^2 \right) \leq c(2)(1 - x_1)^2 M_2^{(n)},$$

where $c(2p)$ and $c(2)$ are some numerical constants and $M_k^{(n)}$ denotes the k -th moment of $A^{(n)}$. We conclude from Hölder's inequality that

$$|\langle \eta^{(n)}, f \rangle - \langle \eta_{\mathbf{x}}^{(n)}, f \rangle| \leq c \|f'\| (1 - x_1) \sqrt{M_2^{(n)} (1 + M_{2p}^{(n)})}.$$

Recall that the sequence $n \mapsto \Phi^{(n)}(q)$ increases for every $q > 0$, so we deduce from Lemma 3.1 that $\sup_n \sqrt{M_2^{(n)} (1 + M_{2p}^{(n)})} < \infty$. Since the dislocation measure ν fulfills the integral condition (2.1), we can complete the proof by dominated convergence. \square

We can now proceed to the proof of Theorem 2.1.

Proof of Theorem 2.1: We first suppose that f and f' are bounded. Then (2.3) follows from Proposition 3.2 and Lemma 3.4 by dominated convergence. Next, we only assume that f is of class \mathcal{C}^1 with $f'(y) = O(y^p)$ as $y \rightarrow \infty$. Then it is easy to construct a sequence $(f_n : n \in \mathbb{N})$ of functions of class \mathcal{C}^1 with f_n and f'_n bounded such that $f_n \rightarrow f$ and $f'_n \rightarrow f'$ pointwise and $\sup_n \|f'_n\| < \infty$ in the sense of Lemma 3.3. This implies that we also have $\sup_n \sup_{x \geq 0} |f_n(x)/(1 + x^{p+1})| < \infty$.

Since all moments of A are finite, we deduce from Lemmas 3.3 and 3.4(i) that

$$\langle \eta, f_n \rangle \rightarrow \langle \eta, f \rangle, \quad \langle \eta, f'_n \rangle \rightarrow \langle \eta, f' \rangle \quad \text{and} \quad \langle \eta_{\mathbf{x}}, f_n \rangle \rightarrow \langle \eta_{\mathbf{x}}, f \rangle$$

where \mathbf{x} is an arbitrary mass-partition. Since $\sup_n \|f'_n\| < \infty$, Lemma 3.4(ii) enables us to apply dominated convergence, and we conclude that (2.3) holds.

We now turn our attention to uniqueness; we consider an arbitrary solution η' to (2.3) and write M'_k for the k -th moment of η' . We have already observed in the introduction that the first moment M'_1 of η' can be computed in terms of the function Φ defined by (2.4). More generally, specifying (2.3) for $f(x) = x^k$ yields an equation of the form

$$\Phi(k(1 - \alpha) - 1) M'_k = \Psi_k(M'_1, \dots, M'_{k-1})$$

for a certain multinomial function Ψ_k . Hence $M'_k = M_k$ for every $k \in \mathbb{N}$, and since we know from Lemma 3.1 that the moment problem for η is determined, this concludes the proof. \square

3.4. *Proof of Lemma 3.3.* We shall finally establish Lemma 3.3 using the framework of homogeneous fragmentations (i.e. self-similar fragmentations with index of self-similarity $\alpha = 0$) with values in the space $\mathcal{P}_{\mathbb{N}}$ of partitions of \mathbb{N} . Given a measure ν which fulfills (2.1), we first consider a homogeneous fragmentation $\Pi = (\Pi(t), t \geq 0)$ with dislocation measure ν and no erosion. For every $t \geq 0$ the random partition $\Pi(t)$ is exchangeable and we write $|\Pi_i(t)|$ for the asymptotic frequency of the i -th block of $\Pi(t)$. The self-similar fragmentation \mathbf{X} is related to Π by a sort of time-change described in Theorem 3.3 of Bertoin (2006), and if we introduce

$$\mathcal{A} = \int_0^\infty \left(\sum_{i=1}^\infty |\Pi_i(t)|^{1-\alpha} \right) dt,$$

then the connexion between Π and \mathbf{X} implies that \mathcal{A} and A have the same distribution.

It will be convenient to approximate \mathcal{A} by Riemann sums. More precisely, for every integer $k \geq 1$ we define

$$\mathcal{A}_k = \frac{1}{k} \sum_{\ell=1}^{k^2} \sum_{i=1}^\infty |\Pi_i(\ell/k)|^{1-\alpha}.$$

The L^1 distance between these two quantities is easily computed in terms of the function Φ defined in (2.4).

Lemma 3.5. *For every $k \geq 1$, we have*

$$\mathbb{E}(|\mathcal{A} - \mathcal{A}_k|) = \frac{1}{\Phi(-\alpha)} - k^{-1} \frac{1 - \exp(-k\Phi(-\alpha))}{\exp(\Phi(-\alpha)/k) - 1}.$$

Proof: The process $|\Pi(\cdot)|^\downarrow$ of the ranked sequence of the asymptotic frequencies of $\Pi(\cdot)$ is a mass-fragmentation; as a consequence $t \mapsto \sum_{i=1}^\infty |\Pi_i(t)|^{1-\alpha}$ decreases and we have $\mathcal{A}_k \leq \mathcal{A}$. Hence $\mathbb{E}(|\mathcal{A} - \mathcal{A}_k|) = \mathbb{E}(\mathcal{A}) - \mathbb{E}(\mathcal{A}_k)$ and the stated formula now follows from the fact that

$$\mathbb{E} \left(\sum_{i=1}^\infty |\Pi_i(t)|^{1-\alpha} \right) = \exp(-t\Phi(-\alpha));$$

see Corollary 2.4(i) and Theorem 3.2 in Bertoin (2006). \square

Next, for every integer n , we write $\Pi^{(n)}$ for the homogeneous fragmentation with dislocation measure $\nu^{(n)}(d\mathbf{x}) = \mathbf{1}_{\{1-x_1 > 1/n\}} \nu(d\mathbf{x})$ and no erosion. We point at the following weak convergence.

Lemma 3.6. *The sequence of processes $(\Pi^{(n)} : n \in \mathbb{N})$ converges in the sense of finite dimensional distributions to Π as $n \rightarrow \infty$.*

Proof: If $\varphi \in \mathcal{P}_{\mathbb{N}}$ is a partition and $k \geq 1$ an integer, then we denote by $\varphi_{|[k]}$ the restriction of φ to the set of the k first integers, $[k] = \{1, \dots, k\}$. We also endow $\mathcal{P}_{\mathbb{N}}$ with the ultra-metric

$$d(\varphi, \varphi') = 1/\sup\{k \geq 1 : \varphi_{|[k]} = \varphi'_{|[k]}\};$$

cf. Lemma 2.6 in [Bertoin \(2006\)](#).

The restriction $\Pi_{|[k]}(t)$ of $\Pi(t)$ to $[k]$ is a Markov chain in continuous time, and we have to verify that $\Pi_{|[k]}^{(n)}$ converges in the sense of finite dimensional distributions to $\Pi_{|[k]}$ as $n \rightarrow \infty$, for each k . This is equivalent to checking the convergence of the corresponding jump rates of the Markov chains.

For every non-trivial partition γ of $[k]$, we write

$$q_\gamma = \lim_{t \rightarrow 0} t^{-1} \mathbb{P}(\Pi_{|[k]}(t) = \gamma).$$

Recall from Theorem 3.1 and Proposition 3.2 in [Bertoin \(2006\)](#) that the jump rate q_γ can be expressed in terms of the dislocation measure ν as

$$q_\gamma = \int_{\mathcal{P}_m} \varrho_{\mathbf{x}}(\gamma) \nu(d\mathbf{x})$$

where $\varrho_{\mathbf{x}}(\gamma)$ is the distribution of Kingman's paintbox process $\pi_{\mathbf{x}}$ based on \mathbf{x} . This means that we consider a sequence ξ_1, \dots, ξ_k of i.i.d. variables with $\mathbb{P}(\xi_1 = i) = x_i$ for $i \geq 1$ and $\mathbb{P}(\xi_1 = 0) = 1 - \sum_{i=1}^{\infty} x_i$, and $\pi_{\mathbf{x}}$ is the exchangeable random partition which is obtained by declaring that two integers $i \neq j$ are in the same block of $\pi_{\mathbf{x}}$ if and only if $\xi_i = \xi_j \neq 0$.

Writing $q_\gamma^{(n)}$ for the jump rate of $\Pi^{(n)}$, we thus have

$$q_\gamma - q_\gamma^{(n)} = \int_{\mathcal{P}_m} \mathbf{1}_{\{1-x_1 \leq 1/n\}} \varrho_{\mathbf{x}}(\gamma) \nu(d\mathbf{x}).$$

It is plain from the paintbox construction that for every mass-partition \mathbf{x} , the probability that the paintbox process based on \mathbf{x} yields the trivial partition on $[k]$ is at least x_1^k . Thus $\varrho_{\mathbf{x}}(\gamma) \leq 1 - x_1^k \leq k(1 - x_1)$ for every non-trivial partition γ of $[k]$. We conclude from (2.1) that $\lim_{n \rightarrow \infty} q_\gamma^{(n)} = q_\gamma$, which establishes our claim. \square

We are now in shape to prove Lemma 3.3

Proof of Lemma 3.3: The space \mathcal{P}_m of mass-partitions is a compact metric space when endowed with the uniform distance (Proposition 2.1 in [Bertoin \(2006\)](#)) and the map $\mathbf{x} \rightarrow \sum_{i=1}^{\infty} x_i^{1-\alpha}$ is continuous and bounded. We write $|\Pi(t)|^\downarrow$ for the sequence of the asymptotic frequency of $\Pi(t)$ ranked in the decreasing order. Recall from Proposition 2.9 in [Bertoin \(2006\)](#) that Lemma 3.6 entails the convergence in the sense of finite dimensional distributions of $(|\Pi^{(n)}(t)|^\downarrow, t \geq 0)$ towards $(|\Pi(t)|^\downarrow, t \geq 0)$. Therefore, for every $k \geq 1$, we have also in the obvious notation

$$\lim_{n \rightarrow \infty} \mathcal{A}_k^{(n)} = \mathcal{A}_k \quad \text{in law.} \tag{3.1}$$

Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a bounded function which is globally Lipschitz-continuous, so by the triangle inequality

$$|\mathbb{E}(f(\mathcal{A})) - \mathbb{E}(f(\mathcal{A}^{(n)}))| \leq |\mathbb{E}(f(\mathcal{A}_k)) - \mathbb{E}(f(\mathcal{A}_k^{(n)}))| + c_f (\mathbb{E}(|\mathcal{A} - \mathcal{A}_k|) + \mathbb{E}(|\mathcal{A}^{(n)} - \mathcal{A}_k^{(n)}|)).$$

It follows readily from Lemma 3.5 that

$$\lim_{k \rightarrow \infty} \mathbb{E}(|\mathcal{A}^{(n)} - \mathcal{A}_k^{(n)}|) = 0 \quad \text{uniformly in } n,$$

so for every $\varepsilon > 0$, we can find an integer k sufficiently large such that

$$\mathbb{E}(|\mathcal{A} - \mathcal{A}_k|) + \mathbb{E}(|\mathcal{A}^{(n)} - \mathcal{A}_k^{(n)}|) \leq \varepsilon/2c_f \quad \text{for all } n,$$

and then we use (3.1) to find an integer n_ε such that $|\mathbb{E}(f(\mathcal{A}_k)) - \mathbb{E}(f(\mathcal{A}_k^{(n)}))| \leq \varepsilon/2$ whenever $n \geq n_\varepsilon$. \square

References

- D. Aldous. Probability distributions on cladograms. In *Random discrete structures (Minneapolis, MN, 1993)*, volume 76 of *IMA Vol. Math. Appl.*, pages 1–18. Springer, New York (1996). [MR1395604](#).
- D. Aldous and A. Bandyopadhyay. A survey of max-type recursive distributional equations. *Ann. Appl. Probab.* **15** (2), 1047–1110 (2005). [MR2134098](#).
- J. Bertoin. Self-similar fragmentations. *Ann. Inst. H. Poincaré Probab. Statist.* **38** (3), 319–340 (2002). [MR1899456](#).
- J. Bertoin. The asymptotic behavior of fragmentation processes. *J. Eur. Math. Soc. (JEMS)* **5** (4), 395–416 (2003). [MR2017852](#).
- J. Bertoin. *Random fragmentation and coagulation processes*, volume 102 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge (2006). [MR2253162](#).
- J. Bertoin and M. Yor. Exponential functionals of Lévy processes. *Probab. Surv.* **2**, 191–212 (electronic) (2005). [MR2178044](#).
- A. F. Filippov. Über das Verteilungsgesetz der Grössen der Teilchen bei Zerstückelung. *Teor. Veroyatnost. i Primenen.* **6**, 299–318 (1961). [MR0140159](#).
- P. Flajolet, P. Poblete and A. Viola. On the analysis of linear probing hashing. *Algorithmica* **22** (4), 490–515 (1998). [MR1701625](#).
- B. Haas and G. Miermont. The genealogy of self-similar fragmentations with negative index as a continuum random tree. *Electron. J. Probab.* **9**, no. 4, 57–97 (electronic) (2004). [MR2041829](#).
- B. Haas, G. Miermont, J. Pitman and M. Winkel. Continuum tree asymptotics of discrete fragmentations and applications to phylogenetic models. *Ann. Probab.* **36** (5), 1790–1837 (2008). [MR2440924](#).
- S. Janson. Brownian excursion area, Wright’s constants in graph enumeration, and other Brownian areas. *Probab. Surv.* **4**, 80–145 (electronic) (2007). [MR2318402](#).
- G. Louchard. The Brownian excursion area: a numerical analysis. *Comput. Math. Appl.* **10** (6), 413–417 (1985) (1984a). [MR783514](#).
- G. Louchard. Kac’s formula, Levy’s local time and Brownian excursion. *J. Appl. Probab.* **21** (3), 479–499 (1984b). [MR752014](#).
- M. Perman and J. A. Wellner. On the distribution of Brownian areas. *Ann. Appl. Probab.* **6** (4), 1091–1111 (1996). [MR1422979](#).
- J. Spencer. Enumerating graphs and Brownian motion. *Comm. Pure Appl. Math.* **50** (3), 291–294 (1997). [MR1431811](#).
- L. Takács. A Bernoulli excursion and its various applications. *Adv. in Appl. Probab.* **23** (3), 557–585 (1991a). [MR1122875](#).
- L. Takács. Conditional limit theorems for branching processes. *J. Appl. Math. Stochastic Anal.* **4** (4), 263–292 (1991b). [MR1136428](#).
- L. Takács. Random walk processes and their applications in order statistics. *Ann. Appl. Probab.* **2** (2), 435–459 (1992). [MR1161061](#).
- L. Takács. On the total heights of random rooted binary trees. *J. Combin. Theory Ser. B* **61** (2), 155–166 (1994). [MR1280604](#).